COPYRIGHT NOTICE

FedUni ResearchOnline
<http://researchonline.ballarat.edu.au>

This is the submitted for peer-review version of the following article:

Anh, L. Q., Kruger, A., Thao, H. (2014). On Hölder calmness of solution mappings in parametric equilibrium problems. TOP, 22(1).331-342

Which has been published in final form at:
<http://doi.org/10.1007/s11750-012-0259-3>

© 2014, Springer.

This is the author's version of the work. It is posted here with permission of the publisher for your personal use. No further distribution is permitted.

On Hölder calmness of solution mappings in parametric equilibrium problems

L.Q. Anh[†], A.Y. Kruger[‡], N.H. Thao[‡]

[†]*Department of Mathematics, Teacher College, Cantho University, Cantho City, Vietnam*

[‡]*School of Science, IT and Engineering, University of Ballarat, Ballarat, Vic, Australia*

Abstract We consider parametric equilibrium problems in metric spaces. Sufficient conditions for the Hölder calmness of solutions are established. We also study the Hölder well-posedness for equilibrium problems in metric spaces.

Keywords Equilibrium problem · calmness · well-posedness

1 Introduction

Optimization is one of the most fertile areas of mathematics. Its conclusions and recommendations play a very important role in both theoretical and applied mathematics. Equilibrium problems were first considered in Blum and Oettli (1994) and since then have been studied by many researchers all over the world. The equilibrium problem model incorporates many other important problems in optimization and other areas such as: variational inequalities, fixed point problems, complementarity, etc. There have been many studies of existence of solu-

Email addresses: quocanh@ctu.edu.vn (L.Q. Anh),
a.kruger@ballarat.edu.au (A.Y. Kruger),
hieuthaonguyen@students.ballarat.edu.au (N.H. Thao).

tions to equilibrium problems (see Sadequi and Alizadeh (2011), Bazán (2001), Bianchi and Schaible (1996), Hai and Khanh (2007a, 2007b), Hai et al. (2009)) and their stability, e.g., semi-continuity in the sense of Berge and Hausdorff (see Anh and Khanh (2004, 2007a, 2007b, 2010), Huang et al. (2006), Khanh and Luu (2007)) or Hölder (Lipschitzian) continuity (see Anh and Khanh (2006, 2008b, 2009), Bianchi and Pini (2003), Li et al. (2009), Li and Li (2011a, 2011b), Mansour and Riahi (2005)).

This paper extends Anh et al. (2011) and studies (l, α) -Hölder calmness of solutions to parametric equilibrium problems. When $\alpha = 1$, this is a kind of calmness property which is in general stronger than the property of the same name usually used in variational analysis. Calmness property of multi-valued mappings has been examined by many authors (see Cánovas et al. (2009), Chuong et al. (2011), Henrion et al. (2002), Ioffe and Outrata (2008), Levy (2000), Ng and Zheng (2009)) in which subdifferentials and coderivatives play the main role. As applications we investigate conditions for Hölder calmness of solutions to optimization problems and well-posedness in the Hölder sense. The last subject is intimately related to the stability property and plays a very important role in studying optimization and variational problems.

The structure of the paper is as follows. Section 2 presents the equilibrium problem model and materials used in the rest of this paper. We establish in Section 3 a sufficient condition for the Hölder calmness of the solution mapping to parametric equilibrium problems. The Hölder well-posedness of equilibrium problems is studied in Section 4.

Throughout the paper, if not explicitly stated otherwise, X, Λ, M are metric spaces and $\mathbb{R}$ is the set of all real numbers while $\mathbb{R}_+$ is the set of all positive numbers. We use $d(\cdot, \cdot)$ for all metrics.

2 Preliminaries

Given a subset $K \subseteq X$ and a function $f : X \times X \rightarrow \mathbb{R}$, a standard *equilibrium problem* is defined as follows:

(EP) Find $\bar{x} \in K$ such that $f(\bar{x}, y) \geq 0$ for all $y \in K$.

The set of solutions to this problem is denoted by S .

In this paper, we consider several extensions of (EP).

The constraint set K and objective function f can be perturbed by parameters $\lambda \in \Lambda$ and $\mu \in M$, respectively. Given a multi-valued mapping $K : \Lambda \rightrightarrows X$, a function $f : X \times X \times M \rightarrow \mathbb{R}$, and a pair $(\lambda, \mu) \in \Lambda \times M$, one can consider a parameterized equilibrium problem:

(EP) $_{\lambda, \mu}$ Find $\bar{x} \in K(\lambda)$ such that $f(\bar{x}, y, \mu) \geq 0$ for all $y \in K(\lambda)$.

The set of solutions to problem (EP) $_{\lambda, \mu}$ is denoted by $S(\lambda, \mu)$.

The approximate version of this problem can be of interest: for each $(\lambda, \mu) \in \Lambda \times M$ and $\varepsilon > 0$,

(EP) $_{\varepsilon, \lambda, \mu}$ Find $\bar{x} \in K(\lambda)$ such that $f(\bar{x}, y, \mu) + \varepsilon \geq 0$ for all $y \in K(\lambda)$.

We denote by $\tilde{S}(\varepsilon, \lambda, \mu)$ the solution set of (EP) $_{\varepsilon, \lambda, \mu}$.

Definition 2.1 For a function $f : X \rightarrow \mathbb{R}$ and positive numbers l, α ,

(i) f is (l, α) -Hölder continuous on a subset $U \subset X$ if

$$|f(x_1) - f(x_2)| \leq ld^\alpha(x_1, x_2) \text{ for all } x_1, x_2 \in U;$$

(ii) f is (l, α) -Hölder calm at $\bar{x}$ on a neighborhood U of $\bar{x}$ if

$$|f(x) - f(\bar{x})| \leq ld^\alpha(x, \bar{x}) \text{ for all } x \in U.$$

We say that f satisfies a certain property on a subset $A \subseteq X$ if it is satisfied at every point of A .

From this definition, it is obvious that Hölder continuity is stronger than Hölder calmness.

To define extensions of these properties for multi-valued mappings we recall the definitions of point-to-set and set-to-set distances. For subsets A, B of X and a point $a \in X$,

$$\begin{aligned} d(a, B) &= \inf_{b \in B} d(a, b); \\ H^*(A, B) &= \sup_{a \in A} d(a, B); \\ H(A, B) &= \max\{H^*(A, B), H^*(B, A)\}; \\ \rho(A, B) &= \sup_{a \in A, b \in B} d(a, b). \end{aligned}$$

Note that H and ρ can take infinite values (if A or B is unbounded). It is also obvious that $H(A, B) \leq \rho(A, B)$ for any subsets A and B , and the inequality can be strict.

Definition 2.2 For a multi-valued mapping $K : \Lambda \rightrightarrows X$ and positive numbers l, α ,

(i) K is (l, α) -Hölder continuous on a subset $U \subset X$ if

$$H(K(\lambda_1), K(\lambda_2)) \leq l d^\alpha(\lambda_1, \lambda_2) \text{ for all } \lambda_1, \lambda_2 \in U;$$

(ii) K is (l, α) -Hölder calm at $\bar{\lambda}$ on a neighborhood U of $\bar{\lambda}$ if

$$H(K(\lambda), K(\bar{\lambda})) \leq l d^\alpha(\lambda, \bar{\lambda}) \text{ for all } \lambda \in U. \quad (1)$$

We will also consider the versions of the properties in Definition 2.2 with H replaced by ρ . In this case, we will talk about the corresponding properties *with respect to ρ* .

Remark 2.1 The calmness in the above definition (when $\alpha = 1$) is a stronger property than the one usually considered in variational analysis. The latter corresponds to replacing H in (1) by H^* (see, e.g., Rockafellar and Wets (1998)). Respectively, (l, α) -calmness is stronger than “calmness $[\alpha]$ ” in Kummer (2009).

We next define *uniform Hölder calmness* as the natural counterpart of the relative Hölder continuity in Anh and Khanh (2007b).

Definition 2.3 For positive numbers m, β, θ , a function $f : X \times X \times M \rightarrow \mathbb{R}$ is (m, β) -Hölder calm at $\bar{\mu}$ on a neighborhood V of $\bar{\mu}$, θ -uniformly over a subset $S \subseteq X$ if

$$|f(x, y, \bar{\mu}) - f(x, y, \mu)| \leq m d^\beta(\bar{\mu}, \mu) d^\theta(x, y), \quad \forall \mu \in V, \forall x, y \in S, x \neq y.$$

If $\theta = 0$, we say that f is (m, β) -Hölder calm at $\bar{\mu}$ on V , uniformly over S .

We next discuss several monotonicity properties some of which are going to play a crucial role in examining the Hölder calmness of the solution mapping of the equilibrium problems $(\text{EP})_{\lambda, \mu}$.

Given a function $f : X \times X \rightarrow \mathbb{R}$, positive numbers h, β , and a subset $S \subseteq X$, consider the following properties.

(M_1) For all $x, y \in S, x \neq y$,

$$f(x, y) + f(y, x) + h d^\beta(x, y) \leq 0. \quad (2)$$

(M_2) For all $x, y \in S$,

$$h d^\beta(x, y) \leq d(f(x, y), \mathbb{R}_+) + d(f(y, x), \mathbb{R}_+). \quad (3)$$

(M_3) For all $x, y \in S, x \neq y$,

$$[f(x, y) \geq 0 \Rightarrow f(y, x) + h d^\beta(x, y) \leq 0].$$

(M_4) For all $x, y \in S, x \neq y$,

$$[f(x, y) < 0 \Rightarrow f(y, x) \geq 0].$$

If any of the above properties is fulfilled, we say that f satisfies the corresponding condition on S with constants h and β (if applicable).

Remark 2.2 Properties (M_1), (M_3) and (M_4) were considered in Anh and Khanh (2006, 2007b, 2008a) where they were called *Hölder strong monotonicity*, *Hölder strong pseudo-monotonicity* and *quasi-monotonicity*, respectively. Property (M_2) is a particular case of the corresponding monotonicity property introduced by Anh and Khanh (see Anh and Khanh (2007b)) for multi-valued mappings. This property has been employed to investigate the Hölder continuity of solution mappings in many articles (see Anh and Khanh (2008a), Li and Li (2011b), Anh et al. (2011).)

The next proposition gives the relationships between these monotonicity properties.

Proposition 2.1

$$(i) \quad (M_1) \Rightarrow (M_2) \Rightarrow (M_3);$$

$$(ii) \quad [(M_3) \& (M_4)] \Rightarrow (M_2).$$

Proof. The following simple observation is used in the proof:

$$d(a, \mathbb{R}_+) = \max\{-a, 0\} \geq -a.$$

(M_1) $\Rightarrow$ (M_2). If (2) holds for some $x \neq y$, then

$$hd^\beta(x, y) \leq -f(x, y) - f(y, x) \leq d(f(x, y), \mathbb{R}_+) + d(f(y, x), \mathbb{R}_+),$$

i.e., (3) holds. When $x = y$, (3) holds automatically.

$(M_2) \Rightarrow (M_3)$. If (3) holds for some $x \neq y$ and $f(x, y) \geq 0$, then $d(f(x, y), \mathbb{R}_+) = 0$ and (3) takes the form

$$hd^\beta(x, y) \leq d(f(y, x), \mathbb{R}_+).$$

It follows from the last inequality that $d(f(y, x), \mathbb{R}_+) > 0$ and consequently $d(f(y, x), \mathbb{R}_+) = -f(y, x)$. Hence, (M_3) holds true.

$[(M_3) \& (M_4)] \Rightarrow (M_2)$. Let (M_3) and (M_4) hold true. We only need to prove (3) when $x \neq y$. If $f(x, y) \geq 0$, then $d(f(x, y), \mathbb{R}_+) = 0$ and (M_3) implies

$$0 < hd^\beta(x, y) \leq -f(y, x) = d(f(y, x), \mathbb{R}_+).$$

Hence, (3) is true. If $f(x, y) < 0$, then (M_4) implies $f(y, x) \geq 0$, and we can apply (M_3) again to show that

$$0 < hd^\beta(x, y) \leq -f(x, y) = d(f(x, y), \mathbb{R}_+).$$

Taking into account that $d(f(y, x), \mathbb{R}_+) = 0$, we conclude that (3) is true in this case too. $\square$

We now give examples showing that implications in Proposition 2.1 can be strict.

Example 2.1 The function $f : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x, y) = x - y$ satisfies (M_2) with $h = \beta = 1$. Indeed,

$$d(f(x, y), \mathbb{R}_+) + d(f(y, x), \mathbb{R}_+) = d(x - y, \mathbb{R}_+) + d(y - x, \mathbb{R}_+) = |x - y| = d(x, y).$$

At the same time, $f(x, y) + f(y, x) = 0$ and (2) is violated for any $x \neq y$. f does not satisfy (M_1) . It is also obvious that f satisfies both (M_3) and (M_4) .

Example 2.2 The function $f : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ defined by $f(x, y) = -\frac{1}{4}(|x|^{\frac{1}{2}} + |y|^{\frac{1}{2}})$ satisfies (M_3) with $h = \sqrt{2}$ and $\beta = \frac{1}{2}$ as $f(x, y) \geq 0$ if and only if $x = y = 0$, it does not satisfy (M_2) . Indeed, for any $y = -x \neq 0$, we have

$$d(f(x, y), \mathbb{R}_+) + d(f(y, x), \mathbb{R}_+) = \frac{1}{2}(|x|^{\frac{1}{2}} + |y|^{\frac{1}{2}}) = |x|^{\frac{1}{2}} < 2|x|^{\frac{1}{2}} = \sqrt{2}d^{\frac{1}{2}}(x, y).$$

We can see that the combination of (M_3) and (M_4) implies (M_2) , but they are not equivalent by considering the function $f(x, y) = -(|x| + |y|)$. This function satisfies (M_2) with $h = \beta = 1$, but breaks (M_4) .

3 The Hölder calmness of the solution mapping

The next theorem gives a sufficient condition for the Hölder calmness of the solution mapping of the problem $(EP)_{\lambda, \mu}$. It improves Theorem 2.1 in Anh et al. (2011). We always assume that solution sets $S(\lambda, \mu)$ are nonempty for all (λ, μ) in a neighborhood of the considered point $(\bar{\lambda}, \bar{\mu})$.

Theorem 3.1 *Consider equilibrium problem $(EP)_{\lambda, \mu}$ and suppose the following conditions hold.*

- (i) *There exist neighborhoods $U(\bar{\lambda})$ of $\bar{\lambda}$ and $V(\bar{\mu})$ of $\bar{\mu}$ and positive numbers n_1, δ_1 and θ such that f is (n_1, δ_1) -Hölder calm at $\bar{\mu}$ on $V(\bar{\mu})$, θ -uniformly over $K(U(\bar{\lambda}))$.*
- (ii) *There exist positive numbers n_2 and δ_2 such that, for all $x \in K(U(\bar{\lambda}))$ and $\mu \in V(\bar{\mu})$, the function $f(x, \cdot, \mu)$ is (n_2, δ_2) -Hölder continuous on $K(U(\bar{\lambda}))$.*
- (iii) *$f(\cdot, \cdot, \bar{\mu})$ satisfies condition (M_2) on $K(U(\bar{\lambda}))$ with constants $h > 0$ and $\beta > \theta$.*
- (iv) *K is (l, α) -Hölder calm at $\bar{\lambda}$ on $U(\bar{\lambda})$ with some positive l and α .*

Then solutions to $(\text{EP})_{\lambda, \mu}$ satisfy the condition of Hölder calmness with respect to ρ : there exist constants $k_1, k_2 > 0$ such that

$$\rho(S(\bar{\lambda}, \bar{\mu}), S(\lambda, \mu)) \leq k_1 d^{\alpha\delta_2/\beta}(\bar{\lambda}, \lambda) + k_2 d^{\delta_1/(\beta-\theta)}(\bar{\mu}, \mu),$$

for all (λ, μ) in a neighborhood of $(\bar{\lambda}, \bar{\mu})$.

Proof. Take $\lambda \in U(\bar{\lambda})$ and $\mu \in V(\bar{\mu})$.

Step 1. We prove that for each $x(\lambda, \bar{\mu}) \in S(\lambda, \bar{\mu})$ and $x(\lambda, \mu) \in S(\lambda, \mu)$,

$$d_1 := d(x(\lambda, \bar{\mu}), x(\lambda, \mu)) \leq \left(\frac{n_1}{h}\right)^{1/(\beta-\theta)} d^{\delta_1/(\beta-\theta)}(\bar{\mu}, \mu). \quad (4)$$

Suppose $x(\lambda, \bar{\mu}) \neq x(\lambda, \mu)$ (if the equality holds, then (4) holds trivially). Because both $x(\lambda, \bar{\mu})$ and $x(\lambda, \mu)$ belong to $K(\lambda)$ and are solutions of $(\text{EP})_{\lambda, \mu}$, one has

$$f(x(\lambda, \bar{\mu}), x(\lambda, \mu), \bar{\mu}) \geq 0; \quad (5)$$

$$f(x(\lambda, \mu), x(\lambda, \bar{\mu}), \mu) \geq 0. \quad (6)$$

At the same time, (iii) implies

$$d\left(f(x(\lambda, \bar{\mu}), x(\lambda, \mu), \bar{\mu}), \mathbb{R}_+\right) + d\left(f(x(\lambda, \mu), x(\lambda, \bar{\mu}), \mu), \mathbb{R}_+\right) \geq h d_1^\beta.$$

Combining this inequality with (5) and (6), we get

$$d\left(f(x(\lambda, \mu), x(\lambda, \bar{\mu}), \bar{\mu}), f(x(\lambda, \mu), x(\lambda, \bar{\mu}), \mu)\right) \geq h d_1^\beta.$$

Because f is (n_1, δ_1) -Hölder calm at $\bar{\mu}$, θ -uniformly over $K(U(\bar{\lambda}))$ by (i), the above relationship implies

$$n_1 d_1^\theta d^{\delta_1}(\bar{\mu}, \mu) \geq h d_1^\beta.$$

This is equivalent to $d_1^{\beta-\theta} \leq \frac{n_1}{h} d^{\delta_1}(\bar{\mu}, \mu)$ from which we get (4) proved.

Step 2. We prove that for each $x(\bar{\lambda}, \bar{\mu}) \in S(\bar{\lambda}, \bar{\mu})$ and $x(\lambda, \bar{\mu}) \in S(\lambda, \bar{\mu})$,

$$d_2 := d(x(\bar{\lambda}, \bar{\mu}), x(\lambda, \bar{\mu})) \leq \left(\frac{2n_2 l^{\delta_2}}{h}\right)^{1/\beta} d^{\alpha\delta_2/\beta}(\bar{\lambda}, \lambda). \quad (7)$$

Suppose $x(\bar{\lambda}, \bar{\mu}) \neq x(\lambda, \bar{\mu})$. (iv) implies that there exist $\bar{x} \in K(\bar{\lambda})$ and $x \in K(\lambda)$ such that

$$d(x(\bar{\lambda}, \bar{\mu}), x) \leq l d^\alpha(\bar{\lambda}, \lambda); \quad (8)$$

$$d(x(\lambda, \bar{\mu}), \bar{x}) \leq l d^\alpha(\bar{\lambda}, \lambda). \quad (9)$$

We get from the definition of $(EP)_{\lambda, \mu}$,

$$f(x(\bar{\lambda}, \bar{\mu}), \bar{x}, \bar{\mu}) \geq 0; \quad (10)$$

$$f(x(\lambda, \bar{\mu}), x, \bar{\mu}) \geq 0. \quad (11)$$

At the same time, (iii) implies

$$d\left(f(x(\bar{\lambda}, \bar{\mu}), x(\lambda, \bar{\mu}), \bar{\mu}), \mathbb{R}_+\right) + d\left(f(x(\lambda, \bar{\mu}), x(\bar{\lambda}, \bar{\mu}), \bar{\mu}), \mathbb{R}_+\right) \geq h d_2^\beta.$$

Combining this inequality with (10) and (11), we get

$$\begin{aligned} & d\left(f(x(\bar{\lambda}, \bar{\mu}), x(\lambda, \bar{\mu}), \bar{\mu}), f(x(\bar{\lambda}, \bar{\mu}), \bar{x}, \bar{\mu})\right) \\ & + d\left(f(x(\lambda, \bar{\mu}), x(\bar{\lambda}, \bar{\mu}), \bar{\mu}), f(x(\lambda, \bar{\mu}), x, \bar{\mu})\right) \geq h d_2^\beta. \end{aligned}$$

Because f is $(n_2 \delta_2)$ -Hölder continuous with respect to the second component in $K(U(\bar{\lambda}))$ by (ii), the last inequality implies that

$$n_2 d^{\delta_2}(x(\lambda, \bar{\mu}), \bar{x}) + n_2 d^{\delta_2}(x(\bar{\lambda}, \bar{\mu}), x) \geq h d_2^\beta.$$

We combine this with (8) and (9) and get

$$n_2 l^{\delta_2} d^{\alpha \delta_2}(\bar{\lambda}, \lambda) + n_2 l^{\delta_2} d^{\alpha \delta_2}(\bar{\lambda}, \lambda) \geq h d_2^\beta,$$

or equivalently $d_2^\beta \leq \frac{2n_2 l^{\delta_2}}{h} d^{\alpha \delta_2}(\bar{\lambda}, \lambda)$. We have (7) proved.

Step 3. For all $x(\bar{\lambda}, \bar{\mu}) \in S(\bar{\lambda}, \bar{\mu})$ and $x(\lambda, \mu) \in S(\lambda, \mu)$, we always have

$$d(x(\bar{\lambda}, \bar{\mu}), x(\lambda, \mu)) \leq d_1 + d_2.$$

From (4) and (7), by taking $k_1 = \left(\frac{2n_2 l^{\delta_2}}{h}\right)^{1/\beta}$ and $k_2 = \left(\frac{n_1}{h}\right)^{1/(\beta-\theta)}$, we get

$$\rho(S(\bar{\lambda}, \bar{\mu}), S(\lambda, \mu)) \leq k_1 d^{\alpha\delta_2/\beta}(\bar{\lambda}, \lambda) + k_2 d^{\delta_1/(\beta-\theta)}(\bar{\mu}, \mu).$$

Therefore, Theorem 3.1 has been proved. $\square$

By using the technique similar to the one in the proof of Theorem 2.1 in Anh and Khanh (2007b), we can show that, under assumption (iii), the solution to $(EP)_{\bar{\lambda}, \bar{\mu}}$ is unique. However, when $(\lambda, \mu) \neq (\bar{\lambda}, \bar{\mu})$, the solutions to $(EP)_{\lambda, \mu}$ do not have to be unique as demonstrated by the following example.

Example 3.1 Let $X = \mathbb{R}$, $\Lambda \equiv M = [0; 1]$, $K(\lambda) = [0; 1]$, $f(x, y, \lambda) = y - x + \lambda$ for all $\lambda \in \Lambda$, and $\bar{\lambda} = 0$.

Then $|f(x, y, \lambda) - f(x, y, \bar{\lambda})| = |\lambda|$. Hence, f is (1.1)-Hölder calm at $\bar{\lambda}$ uniformly over $[0; 1]$. We have $|f(x, y, \lambda) - f(x, z, \lambda)| = |y - z|$ for all $y, z \in [0; 1]$. So $f(x, \cdot, \lambda)$ is (1.1)-Hölder continuous on $[0; 1]$. Therefore, assumptions (i) and (ii) hold. It is clear that condition (iv) also holds. Assumption (iii) is fulfilled as shown in Example 2.1. Hence, Theorem 3.1 derives the Hölder calmness of $S(\cdot)$ at $\bar{\lambda}$. It is not difficult to check that $S(0) = \{0\}$ and $S(\lambda) = [0, \lambda]$ for all $\lambda \in (0; 1]$.

Normally, to receive a property of solution mappings, the problem's hypotheses are also required at the level corresponding to that property. We can see from the preceding theorem that all the hypotheses are related to Hölder continuity and Hölder calmness, except (iii), which is about monotonicity.

The next example indicates the essential role of assumption (iii) in Theorem 3.1.

Example 3.2 Take $X = \mathbb{R}$, $M \equiv \Lambda = [0; 1]$, $K(\lambda) = [-1; 1]$ for all $\lambda \in [0; 1]$. For each $\lambda \in [0; 1]$, consider the function f defined by $f(x, y, \lambda) = \lambda(x + y)$. Take $\bar{\lambda} = 0$.

We have $|f(x, y, \lambda) - f(x, y, \bar{\lambda})| = |x + y| \cdot |\lambda - \bar{\lambda}| \leq 2|\lambda - \bar{\lambda}|$ for all $x, y \in [-1; 1]$. So f is (2.1)-Hölder calm at $\bar{\lambda}$ on $[0; 1]$ uniformly over $[-1; 1]$. At the same time, $|f(x, y, \lambda) - f(x, z, \lambda)| = |\lambda| \cdot |y - z| \leq |y - z|$ for all $y, z \in [-1; 1]$. This means that $f(x, \cdot, \lambda)$ is (1.1)-Hölder continuous on $[-1; 1]$. Hence, conditions (i) and (ii) are fulfilled.

Condition (iv) is also true straightforwardly. However, we have

$$S(0) = [-1; 1], \quad S(\lambda) = \{1\}, \quad \forall \lambda \in (0; 1].$$

So $\rho(S(\lambda), S(0)) = 2$ for any $\lambda \in (0; 1]$.

Therefore, the solution mapping S is not Hölder calm at $\bar{\mu} = 0$. The reason here is that f breaks condition (M_2) . Indeed,

$$d(f(1, 0, 0), \mathbb{R}_+) + d(f(0, 1, 0), \mathbb{R}_+) = 0 < h|1 - 0|^\beta = h, \quad \forall h, \beta > 0.$$

Condition (M_2) in Theorem 3.1 is indispensable.

Remark 3.1 It follows from Proposition 2.1 that the conclusion of Theorem 3.1 remains true if condition (iii) is replaced by either condition (M_1) or conditions (M_3) and (M_4) .

The next proposition aims to illustrate application of Theorem 3.1. For each $(\lambda, \mu) \in \Lambda \times M$, we consider the minimization problem

(MP) Minimize $f(x, \mu)$ subject to $x \in K(\lambda)$,

where $f : X \times M \rightarrow \mathbb{R}$ and $K : \Lambda \rightrightarrows X$. We denote $S(\lambda, \mu) = \{\bar{x} \in K(\lambda) : f(\bar{x}, \mu) = \min_{x \in K(\lambda)} f(x, \mu)\}$ and assume that $S(\lambda, \mu) \neq \emptyset$ for all (λ, μ) near the considered point $(\bar{\lambda}, \bar{\mu})$.

Proposition 3.2 *Consider (MP) and suppose the following conditions hold.*

- (i) *There exist neighborhoods $V(\bar{\mu})$ of $\bar{\mu}$ and $U(\bar{\lambda})$ of $\bar{\lambda}$ and numbers $n_1 > 0$ and $\delta_1 > 0$ such that f is (n_1, δ_1) -Hölder calm at $\bar{\mu}$ on $V(\bar{\mu})$ uniformly over $K(U(\bar{\lambda}))$, i.e.,*

$$|f(x, \mu) - f(x, \bar{\mu})| \leq n_1 d^{\delta_1}(\mu, \bar{\mu})$$

for all $x \in K(U(\bar{\lambda}))$ and $\mu \in V(\bar{\mu})$.

- (ii) *There exist numbers $n_2 > 0$ and $\delta_2 > 0$ such that f is (n_2, δ_2) -Hölder continuous in x on $K(U(\bar{\lambda}))$ uniformly over $\mu \in V(\bar{\mu})$, i.e.,*

$$|f(x, \mu) - f(y, \mu)| \leq n_2 d^{\delta_2}(x, y) \quad (12)$$

for all $\mu \in V(\bar{\mu})$ and $x, y \in K(U(\bar{\lambda}))$, and (12) holds as an equality when $\mu = \bar{\mu}$.

- (iii) *K is (l, α) -Hölder calm at $\bar{\lambda}$ on $U(\bar{\lambda})$ with some $l > 0$ and $\alpha > 0$.*

Then the mapping S is Hölder calm with respect to ρ , i.e., there exist constants $k_1, k_2 > 0$ such that

$$\rho(S(\bar{\lambda}, \bar{\mu}), S(\lambda, \mu)) \leq k_1 d^\alpha(\bar{\lambda}, \lambda) + k_2 d(\bar{\mu}, \mu) \quad (13)$$

for all (λ, μ) in a neighborhood of $(\bar{\lambda}, \bar{\mu})$.

Proof. We define the function $g : X \times X \times M \rightarrow \mathbb{R}$ as follows

$$g(x, y, \mu) = f(y, \mu) - f(x, \mu).$$

We observe that $\bar{x} \in S(\lambda, \mu)$ if and only if $\bar{x} \in K(\lambda)$ and $g(\bar{x}, y, \mu) \geq 0, \forall y \in K(\lambda)$.

So to prove the proposition, it suffices to check that g satisfies the conditions of Theorem 3.1.

We first check condition (i). For every $\mu \in V(\bar{\mu})$ and $x, y \in K(U(\bar{\lambda}))$ we have

$$\begin{aligned} |g(x, y, \mu) - g(x, y, \bar{\mu})| &= |f(y, \mu) - f(x, \mu) - f(y, \bar{\mu}) + f(x, \bar{\mu})| \\ &\leq |f(x, \mu) - f(x, \bar{\mu})| + |f(y, \mu) - f(y, \bar{\mu})| \leq 2n_1 d^{\delta_1}(\mu, \bar{\mu}). \end{aligned}$$

This means that g is $(2n_1, \delta_1)$ -Hölder calm at $\bar{\mu}$ on $V(\bar{\mu})$ uniformly over $K(U(\bar{\lambda}))$.

We have at the same time

$$|g(x, y, \mu) - g(x, z, \mu)| = |f(y, \mu) - f(z, \mu)| \leq n_2 d^{\delta_2}(y, z),$$

i.e., g is (n_2, δ_2) -Hölder continuous with respect to the second component. So conditions (i) and (ii) in Theorem 3.1 are fulfilled.

We now check condition (iii) in Theorem 3.1. For all $x, y \in K(U(\bar{\lambda}))$, we have

$$\begin{aligned} &d(g(x, y, \bar{\mu}), \mathbb{R}_+) + d(g(y, x, \bar{\mu}), \mathbb{R}_+) \\ &= d(f(y, \bar{\mu}) - f(x, \bar{\mu}), \mathbb{R}_+) + d(f(x, \bar{\mu}) - f(y, \bar{\mu}), \mathbb{R}_+) \\ &= |f(x, \bar{\mu}) - f(y, \bar{\mu})| = n_2 d^{\delta_2}(x, y). \end{aligned}$$

So g satisfies condition (M_2) , and (iii) in Theorem 3.1 is fulfilled. Therefore, it follows from Theorem 3.1 that (13) holds true with some $k_1, k_2 > 0$. $\square$

4 The Hölder well-posedness of equilibrium problems

We will denote by $(\mathcal{EP})$ the family of problems $\{(\text{EP})_{\lambda, \mu} : (\lambda, \mu) \in \Lambda \times M\}$ and extend the concept of Lipschitzian well-posedness for optimization problems introduced in Bednarczuk (2007) to equilibrium problems.

Definition 4.1 $(\mathcal{EP})$ is Hölder well-posed at $(\bar{\lambda}, \bar{\mu})$ if $\tilde{S}(0, \bar{\lambda}, \bar{\mu})$ is a singleton and $\tilde{S}$ is Hölder calm at $(0, \bar{\lambda}, \bar{\mu})$ on a neighborhood of $(0, \bar{\lambda}, \bar{\mu})$.

The next theorem gives a sufficient condition for the Hölder well-posedness of $(\mathcal{EP})$. It improves and modifies Theorem 3.1 in Anh et al. (2011).

Theorem 4.1 *Assume $S(\bar{\lambda}, \bar{\mu}) \neq \emptyset$ and the following conditions hold.*

- (i) *There exist neighborhoods $U(\bar{\lambda})$ of $\bar{\lambda}$ and $V(\bar{\mu})$ of $\bar{\mu}$ and positive numbers n_1, δ_1 and θ such that f is (n_1, δ_1) -Hölder calm at $\bar{\mu}$ on $V(\bar{\mu})$, θ -uniformly over $K(U(\bar{\lambda}))$.*
- (ii) *There exist positive numbers n_2 and δ_2 such that, for all $x \in K(U(\bar{\lambda}))$ and $\mu \in V(\bar{\mu})$, the function $f(x, \cdot, \mu)$ is (n_2, δ_2) -Hölder continuous on $K(U(\bar{\lambda}))$.*
- (iii) *$f(\cdot, \cdot, \bar{\mu})$ satisfies condition (M_2) on $K(U(\bar{\lambda}))$ with constants $h > 0$ and $\beta > \theta$.*
- (iv) *K is (l, α) -Hölder calm at $\bar{\lambda}$ on $U(\bar{\lambda})$ with some positive l and α .*

Then $(\mathcal{EP})$ is Hölder well-posed at $(\bar{\lambda}, \bar{\mu})$.

Proof. Take $N = [0, +\infty) \times M$. For $\eta = (\varepsilon, \mu), \eta' = (\varepsilon', \mu') \in N$, consider a function d_N defined by

$$d_N(\eta, \eta') = \max\{|\varepsilon - \varepsilon'|, d(\mu, \mu')\}.$$

Then, (N, d_N) is a metric space. We define a function $g : X \times X \times N \rightarrow \mathbb{R}$ as follows

$$g(x, y, \eta) = f(x, y, \mu) + \varepsilon.$$

To prove the theorem, it suffices to check that g satisfies the conditions of Theorem 3.1.

Take any neighborhood W of 0 in $[0; 1]$. Then for all $\eta = (\varepsilon, \mu) \in W \times V(\bar{\mu})$, $\bar{\eta} = (0, \bar{\mu})$, and $x, y \in K(U(\bar{\lambda}))$, one has

$$|g(x, y, \eta) - g(x, y, \bar{\eta})| = |f(x, y, \mu) - f(x, y, \bar{\mu}) + \varepsilon|$$

$$\leq \varepsilon + |f(x, y, \mu) - f(x, y, \bar{\mu})| \leq \varepsilon + n_1 d^{\delta_1}(\mu, \bar{\mu})$$

$$\leq \varepsilon^{\delta_1} + n_1 d^{\delta_1}(\mu, \bar{\mu}) \leq 2 \max\{1, n_1\} d_N^{\delta_1}(\eta, \bar{\eta})$$

since $\varepsilon \in V \subseteq [0, 1]$ and the Hölder order $\delta_1 \leq 1$. So g is $(2 \max\{1, n_1\} \cdot \delta_1)$ -Hölder calm at $\bar{\eta}$ on $W \times V(\bar{\mu})$ uniformly over $K(U(\bar{\lambda}))$.

We have at the same time

$$|g(x, y, \eta) - g(x, z, \eta)| = |f(x, y, \mu) - f(x, z, \mu)| \leq n_2 d^{\delta_2}(y, z),$$

or g is $(n_2 \cdot \delta_2)$ -Hölder continuous with respect to the second component on $K(U(\bar{\lambda}))$.

Conditions (i) and (ii) of Theorem 3.1 are fulfilled.

We now check condition (iii) of Theorem 3.1. For all $x, y \in K(U(\bar{\lambda}))$, we get

$$\begin{aligned} d(g(x, y, \bar{\eta}), \mathbb{R}_+) + d(g(y, x, \bar{\eta}), \mathbb{R}_+) \\ = d(f(x, y, \bar{\mu}), \mathbb{R}_+) + d(f(y, x, \bar{\mu}), \mathbb{R}_+) \geq h d^\beta(x, y). \end{aligned}$$

This means that g satisfies condition (iii) of Theorem 3.1 and we have all its hypotheses satisfied. Therefore, the mapping of solutions to $(\mathcal{EP})$ is both Hölder calm and single-valued at $(0, \bar{\eta})$ which combined with Definition 4.1 gives the conclusion of the theorem. $\square$

5 Conclusion

Assuming Hölder calmness and Hölder continuity in Hausdorff distance, we have established the Hölder calm property of the solution mapping with respect to ρ . This obviously implies the Hölder calm property in Hausdorff distance. We have established a sufficient condition for the Hölder well-posedness of equilibrium problems. These may be extended to many other classes of problems.

Acknowledgements

The authors wish to thank Phan Quoc Khanh and the anonymous referees for the careful reading of the paper and valuable comments and suggestions.

References

- Anh LQ, Duy TQ, Thao NH, Van DTM (2011) On the Hölder calm continuity and Hölder well-posedness of parametric equilibrium problems in metric spaces. *Sci J Cantho Uni*, 19b: 70–79 (in Vietnamese)
- Anh LQ, Khanh PQ (2004) Semicontinuity of the solution set of parametric multivalued vector quasiequilibrium problems. *J Math Anal Appl* 294:699–711
- Anh LQ, Khanh PQ (2006) On the Hölder continuity of solutions to multivalued vector equilibrium problems. *J Math Anal Appl* 321:308–315
- Anh LQ, Khanh PQ (2007a) On the stability of the solution sets of general multivalued vector quasiequilibrium problems. *J Optim Theory Appl* 135:271–284
- Anh LQ, Khanh PQ (2007b) Uniqueness and Hölder continuity of the solution to multivalued equilibrium problems in metric spaces. *J Glob Optim* 37:449–465
- Anh LQ, Khanh PQ (2008a) Sensitivity analysis for multivalued quasiequilibrium problems in metric spaces: Hölder continuity of solutions. *J Glob Optim* 42:515–531
- Anh LQ, Khanh PQ (2008b) Semicontinuity of the approximate solution sets of multivalued quasiequilibrium problems. *Numer Funct Anal Optim* 29:24–42

- Anh LQ, Khanh PQ (2009) Hölder continuity of the unique solution to quasiequilibrium problems in metric spaces. *J Optim Theory Appl* 141:37–54
- Anh LQ, Khanh PQ (2010) Continuity of solution maps of parametric quasiequilibrium problems. *J Glob Optim* 46:247–259
- Bazán FF (2001) Existence theorem for generalized noncoercive equilibrium problems. *SIAM J Optim* 11:675–690
- Bednarczuk E (2007) *Stability Analysis for Parametric Vector Optimization Problems*. Polish Academy of Sciences, Warszawa
- Bianchi M, Pini R (2003) A note on stability for parametric equilibrium problems. *Oper Res Lett* 31:445–450
- Bianchi M, Schaible S (1996) Generalized monotone bifunctions and equilibrium problems. *J Optim Theory Appl* 90:31–43
- Blum E, Oettli W (1994) From optimization and variational inequalities to equilibrium problems. *Math Student* 63:123–145
- Cánovas MJ, Dontchev AL, López MA, Parra J (2009) Isolated calmness of solution mappings in convex semi-infinite optimization. *J Math Anal Appl* 350:829–837
- Chuong TD, Kruger AY, Yao JC (2011) Calmness of efficient solution maps in parametric vector optimization. *J Glob Optim* 51:677–688
- Hai NX, Khanh PQ (2007a) Existence of solutions to general quasiequilibrium problems and applications. *J Optim Theory Appl* 133:317–327
- Hai NX, Khanh PQ (2007b) The solution existence of general variational inclusion problems. *J Math Anal Appl* 328:1268–1277

- Hai NX, Khanh PQ, Quan NH (2009) On the existence of solutions to quasivariational inclusion problems. *J Glob Optim* 45:565–581
- Henrion R, Jourani A, Outrata J (2002) On the calmness of a class of multifunctions. *SIAM J Optim* 13:603–618
- Huang NJ, Li J, Thompson HB (2006) Stability for parametric implicit vector equilibrium problems. *Math Comput Modelling* 43:1267–1274
- Ioffe A, Outrata J (2008) On metric and calmness qualification conditions in subdifferential calculus. *Set-Valued Anal* 16:199–227
- Khanh PQ, Luu LM (2007) Lower and upper semicontinuity of the solution sets and the approximate solution sets to parametric multivalued quasivariational inequalities. *J Optim Theory Appl* 133:329–339
- Kummer B (2009) Inclusions in general spaces: Hoelder stability, solution schemes and Ekeland’s principle. *J Math Anal Appl* 358:327–344
- Levy AB (2000) Calm minima in parameterized finite-dimensional optimization. *SIAM J Optim* 11:160–178
- Li XB, Li SJ (2011a) Continuity of approximate solution mapping for parametric equilibrium problems. *J Glob Optim* 51:541–548
- Li SJ, Li XB (2011b) Hölder continuity of solutions to parametric weak generalized Ky Fan inequality. *J Optim Theory Appl* 149:540–553
- Li SJ, Li XB, Teo KL (2009) The Hölder continuity of solutions to generalized vector equilibrium problems. *European J Oper Res* 199:334–338
- Mansour MA, Riahi H (2005) Sensitivity analysis for abstract equilibrium problems. *J Math Anal Appl* 306:684–691

- Ng KF, Zheng XY (2009) Calmness for L-subsmooth multifunctions in Banach spaces. *SIAM J Optim* 19:1648–1673
- Sadequi I, Alizadeh CG (2011) Existence of solutions of generalized vector equilibrium problems in reflexive Banach spaces. *Nonlinear Anal* 74:2226–2234
- Rockafellar RT, Wets RJB (1998) *Variational Analysis*, volume 317 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin.